

Evolutionary trade-off between weapons and testes

Leigh W. Simmons^{*†} and Douglas J. Emlen^{*}

^{*}Centre for Evolutionary Biology, School of Animal Biology (M092), University of Western Australia, Crawley, Western Australia 6009, Australia; and [†]Division of Biological Sciences, University of Montana, Missoula, MT 59812

Edited by Bert Hölldobler, Arizona State University, Tempe, AZ, and approved September 13, 2006 (received for review April 28, 2006)

It has long been recognized that male mating competition is responsible for the evolution of weaponry for mate acquisition. However, when females mate with more than one male, competition between males can continue after mating in the form of sperm competition. Theory predicts that males should increase their investment in sperm production as sperm competition is increased, but it assumes that males face a trade-off between sperm production and other life-history traits such as mate acquisition. Here, we use a genus of horned beetle, *Onthophagus*, to examine the trade-off between investment in testes required for fertilizations and investment in weapons used to obtain matings. In a within-species study, we prevented males from developing horns and found that these males grew larger and invested relatively more in testes growth than did males allowed to grow horns. Among species, there was no general relationship between the relative sizes of horns and testes. However, the allometric slope of horn size on body size was negatively associated with the allometric slope of testes size on body size. We suggest that this reflects meaningful evolutionary changes in the developmental mechanisms regulating trait growth, specifically in the degree of nutrition-dependent phenotypic plasticity versus canalization of traits. Finally, we show how this resource allocation trade-off has influenced the evolutionary diversification of weapons, revealing a rich interplay between developmental trade-offs and both pre- and postmating mechanisms of sexual competition.

secondary sexual traits | sperm competition | testes size | beetle horns

Competition among males for mating opportunities has long been recognized as a potent selective pressure shaping the evolution of exaggerated secondary sexual weapons, such as deer antlers and beetle horns (1). However, when females mate with more than one male, competition between males can continue after copulation so that sperm from different males must compete to fertilize eggs (2). Theory predicts that sperm competition should favor increased male allocation to sperm production (3), and there is accumulating empirical support for the basic prediction of this theory, that sperm competition can favor increased male expenditure on sperm production. Thus, among species from a broad range of taxa, positive evolutionary associations have been found between male investment in gametogenic tissue (testes size) and estimates of the strength of selection generated by sperm competition (4–8).

Sperm competition theory rests on the fundamental assumption that males have a limited resource pool with which to invest in reproduction and that they must trade off resources between gaining matings and fertilizations (3). This underlying assumption of a trade-off between investment in testes and investment in other life-history traits has received less attention from empiricists than have the predicted outcomes of sperm competition theory. Nonetheless, there is some evidence to suggest that energetically expensive traits such as immunity can trade off against testis size and ejaculate quality (9, 10), and across species of fruit flies in the genus *Drosophila*, species developing larger testes require longer periods of sexual maturation (11, 12). Finally, a recent comparative study of bats revealed a trade-off between brain size and testes size, two metabolically expensive organs (13).

But do males face a trade-off between investments in competing for fertilizations and competing for access to mates? Studies of species in which males follow alternative life histories that are associated with alternative mating tactics suggest that they might. For example in salmon, anadromic males invest heavily in body growth and in weapons for competing for spawning females. In contrast parr mature at a small body size and invest more in their testes, producing sperm with greater levels of ATP activity, motility, and fertilization capacity (14, 15).

Beetles in the genus *Onthophagus* are notable for the size and diversity of their horns. Horns can comprise a significant proportion of male body size and can project from any of five locations on the head or thorax. Horns can be expressed in both sexes but in most species are expressed in males only (sexual dimorphism) or in just the largest males (male dimorphism) (16). Males use their horns to block the entrance to breeding tunnels containing females, and males with larger horns are competitively superior (17, 18) and attain a higher reproductive success than their short-horned conspecifics (19). The horns of onthophagines thereby represent typical sexually selected weapons that increase the number of matings obtained.

Sperm competition can also be an important selective pressure in onthophagine mating systems. In male dimorphic species, minor or hornless males adopt the alternative mating tactic of sneaking copulations with females that are guarded by major or horned males (17, 18, 20). In the species that have been studied, a male's fertilization success depends on the amount of sperm inseminated relative to other males (21, 22), an assumption made in sperm competition game theory (3), and the proportion of sneak males in a population is associated with both the relative sizes of the testes, and the numbers of sperm produced (23), suggesting that sperm competition favors large testes size in these beetles. Thus, onthophagine beetles appear to have experienced a history of selection for significant investment into both horns and testes and for this reason are ideal for testing for a developmental allocation trade-off between these traits.

Production of horns is known to come at the expense of the development of other morphological structures (24). Within natural populations of onthophagine beetles, the relative sizes of horns are often negatively correlated with the relative sizes of other traits including eyes, wings, and antennae (24), and within-species perturbation experiments have confirmed that altering allocation to one trait can impact growth of another [e.g., artificial selection for increased relative horn size reduced relative eye size (25), and ablation of genital capsules in larvae resulted in males developing horns at smaller body sizes compared with intact males (26)].

Here we use an experimental manipulation of horn development to provide evidence that male *Onthophagus nigriventris* trade-off investment in the weapons used to obtain matings for investment in testes required for fertilizations. We then conduct

Author contributions: L.W.S. and D.J.E. designed research; L.W.S. and D.J.E. performed research; L.W.S. and D.J.E. analyzed data; and L.W.S. and D.J.E. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS direct submission.

[†]To whom correspondence should be addressed. E-mail: lsimmons@cyllene.uwa.edu.au.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. Evidence for an allocation trade-off between weapons (horns) and testes in the beetle genus *Onthophagus* (Coleoptera: Scarabaeidae). (A) Within the species *O. nigriventris*, large males produce two horns on the thorax. (B) Males engineered to be hornless (\square) enclosed into larger adults with testes that were disproportionately large compared with control males that developed horns (\bullet). The common regression line is shown with its 95% confidence limits. (B Inset) The residual testes weights for cauterized and control beetles after controlling for body weight. (C and D) In a comparative analyses of 25 congeneric species, the steepness of the allometric slope of log testes weight on log body weight declined across species as both body size (C) and the steepness of the scaling relationship between horn length and body size (D) increased. Photograph by O. Helmy.

a comparative analysis across 25 species of *Onthophagus* to determine whether this resource allocation trade-off has influenced the evolutionary diversification of weapons in this group of beetles. Our data reveal a rich interplay between developmental trade-offs and both pre- and postmating mechanisms of sexual competition.

Results

Experimental Manipulation of Horn Development. Male *O. nigriventris* produce a pair of thoracic horns, one of which can reach up to 40% of the male's total body length (Fig. 1A). We experimentally engineered males not to invest in horn growth by cauterizing in late instar larvae the area of proliferating cells that would otherwise become the principal thoracic horn. Cauterized beetles grew larger (112.4 ± 7.0 mg) than uncaterized beetles (87.3 ± 3.9 mg) ($t_{47} = 2.93$, $P = 0.005$), and they failed to produce a thoracic horn (Fig. 1B). An ANCOVA with treatment (cauterized versus control) as the main effect and log body weight as the covariate explained 71% of the variance in log testes weight ($F_{2,45} = 55.53$, $P < 0.0001$). The interaction between the covariate and main effect was not significant ($F_{1,44} = 0.15$, $P = 0.704$) indicating homogeneity of slopes, and was removed from the model before evaluating the independent partial effect of treatment (27). Testes weight increased with body weight ($F_{1,45} = 12.98$, $P = 0.0008$; common slope 0.89 ± 0.14), but cauterized males developed disproportionately larger testes than did control males ($F_{1,45} = 52.84$, $P < 0.0001$) (Fig. 1B).

Comparative Analyses. We looked for evidence of an evolutionary trade-off between investment in horns and testes across 25 species of *Onthophagus* (Table 1, which is published as supporting information on the PNAS web site). If the phenotypic trade-off found in *O. nigriventris* has affected long term patterns of either horn or testes evolution, then we might expect species in which males invest heavily in horn development for mate acquisition to have a lower investment in testes growth.

We first used a general linear model to test whether testes weight was negatively correlated with horn size, including species mean body size as a covariate in the analysis (all variables were log transformed for analysis). The full model explained 92% of the variation in testes weight ($F_{2,22} = 130.47$, $P < 0.001$). Not surprisingly, larger species had larger testes (effect estimate 0.64 ± 0.06 , $F_{1,22} = 119.46$, $P < 0.001$). Contrary to our prediction however, we detected a nonsignificant trend toward a positive relationship between the relative sizes of these primary and secondary sexual structures (effect estimate 0.05 ± 0.02 , $F_{1,22} = 3.31$, $P = 0.083$), suggesting that developmental allocation trade-offs may not have been important in shaping the evolution of relative trait sizes.

We also looked at a second pattern. We used the slopes of the trait-size body size scaling relationships (allometry) as a measure of the species-specific pattern of trait development (28, 29), and we tested for an evolutionary relationship between these allometric slopes. Because among-individual variation in body size in these beetles is largely determined by the nutritional envi-

previous within-species studies which showed that negative phenotypic associations between horns and body structures such as eyes, antennae or wings, increase in magnitude with increasing proximity of the developing horns to adjacent body structures (24). Thus, it might be that sperm competition constrains the origin of novel thoracic horns but not head horns, because thoracic horns are more likely to draw from the same resource pool during development.

In conclusion, our data reveal the trade-off between male investment in gaining matings and fertilizations explicit in evolutionary models of ejaculate expenditure. They suggest that sperm competition can play an important, although not always intuitive, role in shaping the evolution of secondary sexual traits used in conventional contest competition, and thereby contribute to the evolutionary diversification of male morphology.

Methods

Experimental Manipulation of Horn Development. We looked for a trade-off between horn growth and testes growth in *O. nigriventris* by experimentally manipulating horn development. Experimental *O. nigriventris* were derived from adult beetles collected from pastures of the Parker Ranch on the island of Hawaii. Mated females were established in plastic breeding chambers (8 cm diameter \times 20 cm deep) in the laboratory (University of Montana) and maintained in growth chambers at 25°C with unlimited access to cow dung. Females build a series of individual brood balls, each of which provides the resources for the development of a single offspring from hatching until adult emergence. Brood balls were sieved from breeding chambers and placed in shallow soil-filled containers in the same growth chambers. Individual broods were opened and larvae checked for their stage of development. Toward the end of the third larval instar, during the gut purge, the area of proliferating cells that would otherwise become the principal thoracic horn was cauterized by application of a hyphrecator. The larvae were held in place with forceps during cauterization and care was taken not to discolor the larval cuticle. The larvae were then returned to their individual brood balls and allowed to continue their development. Emerging adult beetles were checked for horn growth. No cauterized beetles developed horns. A series of controls were established in which larvae were removed from their broods, handled in a similar manner to cauterized beetles, and then returned to their broods. Seven days after adult emergence (after testes have matured), beetles were weighed and then dissected, and their testes were removed and weighed to an accuracy of 0.01 mg. Weights were log transformed before statistical analyses.

Comparative Analyses. We collected data on thorax width (as a measure of body size), testes weight, body weight, and horn morphology from 25 species of *Onthophagus*. For all species beetles were sampled from the field, collected from fresh animal dung. Beetles were returned to the laboratory and maintained at 25°C with access to fresh dung for 7 days. Before dissection, beetles were held without access to dung or water for 24 h. This procedure was adopted to ensure that beetles were sexually mature, of equivalent recent mating history, and that they had

purged their guts before weight determination. Thorax width was cubed to convert it to the same scale as our weight measures.

The genus exhibits complex patterns of variation in horn morphology, developing horns at one or several of five locations on the head and thorax (16, 32). We used two metrics of horn morphology (Table 1). First we measured the length of the largest or most exaggerated horn and again cubed this value to achieve the same scale as our weight measures. Second, we examined the degree of developmental plasticity in horn growth by calculating the scaling relationship between horn size and body size. For many species of *Onthophagus*, the scaling relationship of horn size on body size deviates from linearity because of the occurrence of minor males that in some species are hornless, whereas in others develop only rudimentary horns (32). We therefore fitted data for each species to the following equation:

$$\text{horn length} = y_0 + \frac{ax^b}{c^b + x^b}, \quad [1]$$

where y_0 is equal to the minimum horn length, a describes the range of horn lengths present in the sample, b the maximum rate of increase in horn size per unit increase in body size, c represents the body size at the point of inflection of the sigmoid, and x represents our measure of body size, thorax width. We used the software package SigmaPlot (SYSTAT) to estimate the parameter b from this equation and used this value as our measure of horn allometry for each species. Where species possessed both head and thoracic horns, we calculated the scaling relationship for the largest or most exaggerated horn.

Again, in addition to measuring a species' absolute testes weight, we quantified each species' degree of developmental plasticity in testes growth as the allometric slope of log testes weight on log body weight. We calculated the major axis slope (MA in Table 1) because we are interested in the true relationship between these two variables, both were measured with error, and both were measured on the same scale (49).

Of the species listed in Table 1, all but *O. blackwoodensis*, *O. nodulifer*, and *O. rupicapra* were included in a recent phylogeny, based on regions of four nuclear and three mitochondrial genes (3,315 bp total; 837 parsimony-informative) from 48 *Onthophagus* species and three outgroups (16, 32). Based on this phylogeny, we used the software package CAIC (33) to calculate phylogenetically independent contrasts. We also used the phylogeny to test for correlated evolutionary changes between gains of novel horns (on the head; on the thorax) and changes in the level of sperm competition as estimated by the presence/absence of hornless minor (sneak) males (male-dimorphism; male-monomorphism; see results for justification). For these analyses, we used the concentrated changes test (50) as implemented in MacClade 4 (36).

We thank everyone who sent beetles from their travels and Quenna Szafran for conducting the cauterizations. The work was funded by grants from the Australian Research Council and the University of Western Australia Small Grant Scheme (to L.W.S.) and by National Science Foundation Grant IOB PECASE 0092873 (to D.J.E.) At the University of Montana, beetles were housed under U.S. Department of Agriculture Animal and Plant Health Inspection Service Permit 45534.

1. Darwin C (1871) *The Descent of Man and Selection in Relation to Sex* (John Murray, London).
2. Parker GA (1970) *Biol Rev* 45:525–567.
3. Parker GA (1998) in *Sperm Competition and Sexual Selection*, eds Birkhead TR, Møller AP (Academic, London), pp 3–54.
4. Stockley P, Gage MJG, Parker GA, Møller AP (1997) *Am Nat* 149:933–954.
5. Hosken DJ (1997) *Proc R Soc London Ser B* 264:385–392.
6. Gage MJG (1994) *Proc R Soc London Ser B* 258:247–254.
7. Harcourt AH, Purvis A, Liles L (1995) *Funct Ecol* 9:468–476.
8. Byrne PG, Roberts JD, Simmons LW (2002) *J Evol Biol* 15:347–355.

9. Hosken DJ (2001) *Curr Biol* 11:379–380.
10. Simmons LW, Roberts B (2005) *Science* 309:2031.
11. Pitnick S (1996) *Am Nat* 148:57–80.
12. Pitnick S, Miller GT (2000) *Heredity* 84:416–426.
13. Pitnick S, Jones KE, Wilkinson GS (2006) *Proc R Soc London Ser B* 273:719–724.
14. Vladic TV, Järvi T (2001) *Proc R Soc London Ser B* 268:2375–2381.
15. Gage MJG, Stockley P, Parker GA (1995) *Philos Trans R Soc London B* 350:391–399.
16. Emlen DJ, Marangelo J, Ball B, Cunningham CW (2005) *Evolution* (Lawrence, KS) 59:1060–1084.

17. Emlen DJ (1997) *Behav Ecol Sociobiol* 41:335–342.
18. Moczek AP, Emlen DJ (2000) *Anim Behav* 59:459–466.
19. Hunt J, Simmons LW (2001) *Proc R Soc London Ser B* 268:2409–2414.
20. Hunt J, Simmons LW (2002) *J Evol Biol* 15:784–795.
21. Tomkins JL, Simmons LW (2000) *Proc R Soc London Ser B* 267:1547–1553.
22. Simmons LW, Beveridge M, Krauss S (2004) *Behav Ecol Sociobiol* 57:164–173.
23. Simmons LW, Tomkins JL, Hunt J (1999) *Proc R Soc London Ser B* 266:145–150.
24. Emlen DJ (2001) *Science* 291:1534–1536.
25. Nijhout HF, Emlen DJ (1998) *Proc Natl Acad Sci USA* 95:3685–3689.
26. Moczek AP, Nijhout HF (2004) *Am Nat* 163:184–191.
27. Engqvist L (2005) *Anim Behav* 70:967–971.
28. Emlen DJ, Nijhout HF (2000) *Annu Rev Entomol* 45:661–708.
29. Emlen DJ, Allen CE (2004) *Integr Comp Biol* 43:617–634.
30. Hunt J, Simmons LW (2000) *Evolution (Lawrence, KS)* 54:936–941.
31. Emlen DJ (1994) *Proc R Soc London Ser B* 256:131–136.
32. Emlen DJ, Hunt J, Simmons LW (2005) *Am Nat* 166:S42–S68.
33. Purvis A, Rambaut A (1995) *Comput Appl Biosci* 11:247–251.
34. Pomfret JC, Knell RJ (2006) *Anim Behav* 71:567–576.
35. Kawano K (2004) *Am Nat* 163:1–15.
36. Maddison WP, Maddison DR (1999) *MacClade 4* (Sinauer, Sunderland, MA).
37. Via S, Lande R (1985) *Evolution (Lawrence, KS)* 39:505–522.
38. Schluter D (1996) *Evolution (Lawrence, KS)* 50:1766–1774.
39. Lande R (1979) *Evolution (Lawrence, KS)* 33:402–416.
40. Messina FJ, Slade AF (1999) *Physiol Entomol* 24:358–363.
41. Messina FJ, Fry JD (2003) *J Evol Biol* 16:501–509.
42. Stern DL, Emlen DJ (1999) *Development (Cambridge, UK)* 126:1091–1101.
43. Cotton S, Fowler K, Pomiankowski A (2004) *Proc R Soc London Ser B* 271:771–783.
44. Simmons LW, Tomkins JL (1996) *Evol Ecol* 10:97–104.
45. Goberdhan DC, Wilson C (2002) *Dev Genes Evol* 212:196–202.
46. Shingleton AW, Das J, Vinicius L, Stern DL (2005) *PLoS Biol* 3:1607.
47. Eberhard WG, Huber BA, Rodriguez RLS, Briceño RD, Salas I, Rodríguez V (1998) *Evolution (Lawrence, KS)* 52:415–431.
48. House CM, Simmons LW (2005) *J Evol Biol* 18:1281–1292.
49. Harvey PH, Pagel MD (1991) *The Comparative Method in Evolutionary Biology* (Oxford Univ Press, Oxford).
50. Maddison WP (1990) *Evolution (Lawrence, KS)* 44:539–557.